RITE CARE RESULTS



RITE CARE WORKS

- √ Low rates of emergency hospital admissions
- √ Low rates of preventable hospitalizations
- √ Improved access to primary care
- √ Fewer pregnant women who smoke
- √ Healthier pregnancies
- √ Fewer infant deaths
- √ Healthier infants and children

RIte Care is a health insurance program that has a 20-year track record of successfully providing high-quality, affordable health care for Rhode Island children and families who would otherwise be uninsured. When Rhode Island children and families are covered efficiently and cost-effectively, we all win.


When it comes to better health outcomes for children and families - RIte Care works!

November 2013

RITE CARE RESULT #1: Low Rates of Emergency Hospital Admissions for Children & Adults


Health care that is delivered in the emergency department is among the most costly types of health care. Access to primary and preventive health care providers reduces emergency room use and costly emergency hospital admissions.

Children and adults under age 65 with Medicaid coverage have low rates of emergency admissions to the hospital. Thirty-nine percent of admissions to the hospital for Medicaid members are emergencies, the same as those privately insured. The percentage of emergency admissions for children and adults who are uninsured has increased 30% over the past decade, from 64% in 2001 to 83% in 2011. (see figure A)




*Medicaid includes fee-for-service and managed care (i.e. Rite Care and Rhody Health Partners)

RITE CARE RESULT #2: BETTER ACCESS TO PRIMARY CARE = FEWER PREVENTABLE HOSPITALIZATIONS



*Rite Care includes managed care and fee-for-service Medicaid



*Medicaid includes fee-for-service and managed care (i.e. RIte Care and Rhody Health Partners)

Access to health insurance and to primary care can reduce preventable hospitalizations for children and adults. Preventable hospitalizations are those that result from conditions that could have been treated through routine primary and preventive care. These conditions include asthma, pneumonia, diabetes, hypertension, chronic obstructive pulmonary disease, etc., all of which can be well-controlled through the regular care of a health care provider.

Children in Rhode Island who have either RIte Care or private health insurance have fewer preventable hospitalizations than uninsured children. The percentage of preventable hospitalizations for children who were uninsured in Rhode Island nearly tripled between 2001 and 2011 (from 7% to 20%), while preventable hospitalizations for children with RIte Care coverage decreased by a third, from 12% to 8%. (see figure B)

Adults with Medicaid coverage also have relatively low rates of preventable hospitalizations. The percentage of preventable hospitalizations for Rhode Island adults who are uninsured (17%) is nearly two times higher than the rate of adults with Medicaid (10%). (see figure C)

Children and adults with health coverage, including Medicaid and RIte Care, have improved access to preventive care, leading to fewer costly and preventable hospitalizations. Children who receive timely immunizations and see their primary care physician throughout childhood have better health outcomes than those who do not. Primary care for adults helps to ensure that chronic diseases are well-managed and that serious illnesses are detected early.¹

RITE CARE RESULT #3: HEALTHIER PREGNANCIES, HEALTHIER CHILDREN

Women who get regular health care before and during pregnancy, who don't smoke while they are pregnant and who have children at least 18 months apart are less likely to have babies with health problems.

Early prenatal care is important to identify and treat health problems. Women receiving late or no prenatal care are at increased risk of poor birth outcomes, such as having babies who are low birthweight or who die within the first year of life.²




figure D

Fewer women with RIte Care coverage smoke during pregnancy. Smoking during pregnancy increases the likelihood of poor birth outcomes, including having a baby that is stillborn, preterm or low birthweight.^{3,4} The percentage of women enrolled in RIte Care who smoked while pregnant decreased by 53% between 1993 and 2011, from 32% to 15%. (see figure D)





figure E

Better access to health care means that more women are able to reduce the risks associated with short interval births. Women who have short interval births (less than 18 months between one child and the next) face more health risks to the mother and the infant. The percentage of women with RIte Care coverage who have short interval births declined by 41% (from 41% to 24%) between 1993 and 2011. This marks the first time since 2008 that the RIte Care rate is lower than those with private insurance (26%).



RITE CARE RESULT #4: FEWER INFANT DEATHS AND HEALTHIER BABIES

The infant mortality rate is an important measure of the well-being of infants, children and pregnant women. Infant mortality is related to a variety of factors, including women's health status, quality and access to medical care, socio-economic conditions and public health practices.⁶





figure F



Fewer Rhode Island babies are dying in the first year of life. Rhode Island's infant mortality rate for publicly-insured infants dropped from 9.8 infant deaths per 1,000 live births in 1993 to 7.7 in 2011. (see figure F) However, there is more work to be done in this area.

In 2010, the most recent year for which national comparison data are available, Rhode Island's overall infant mortality rate (for all infants regardless of type of health insurance coverage) of 7.1 per 1,000 live births was above the record low U.S. rate of 6.1 death per 1,000 live births. This marks the first time since 2007 that Rhode Island's overall infant mortality rate was above the national average. From 2009 to 2010, Rhode Island slipped from 24th best in the U.S. and 5th best in New England to 37th best in the U.S. and to last in New England on this measure.^{7,8}

An infant's birthweight is a key indicator of newborn health. Children born at low birthweight face greater risks of physical and developmental health problems and death than infants of normal birthweight. They also are more likely to have long-term disabilities and exhibit poor school performance. 9,10,11

National trends showed overall increases in the percentage of infants with low birthweight (<2,500 grams or 5 pounds, 8 ounces) between 1990 and 2006, with small declines since then. Rhode Island mirrored that trend, with RIte Care low birthweight rates peaking at 9.6% in 2003, and then improving to 8.4% in 2011. (see figure G) Rhode Island's overall low birthweight rate was consistently below the national average between 2004 and 2011. ^{12,13}

References

- ¹ Institute of Medicine. (2009). America's uninsured crisis: Consequences for health and health care. Washington, DC: The National Academies Press.
- ² Child Trends Data Bank. (2012). Late or no prenatal care. Retrieved October 16, 2013, from www.childtrendsdatabank.org
- 3.10 U.S. Department of Health and Human Services. (2010). How tobacco smoke causes disease: The biology and behavioral basis for smoking-attributable disease: A report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services.
- ⁴U.S. Department of Health and Human Services, Office on Women's Health. (2009). *Prenatal care frequently asked questions*. Retrieved October 16, 2013, from www.womenshealth.gov/publications/our-publications/fact-sheet/prenatal-care.cfm
- ⁵Rhode Island Department of Health. (2012). 2013-2015 Rhode Island preconception health strategic plan. Retrieved October 16, 2013, from www.health.ri.gov/publications/strategicplans/2013RhodeIslandPreconceptionHealth2015.pdf
- 6.11 Federal Interagency Forum on Child and Family Statistics. (2012). America's children: Key national indicators of well-being 2012. Washington, DC: Government Printing Office.
- Murphy, S., et al. (2013). Deaths: Final Data for 2010. National Vital Statistics Reports, 61(4). Hyattsville, MD: Centers for Disease Control and Prevention.
- 8,13 The Annie E. Casey Foundation, KIDS COUNT Data Center, datacenter.kidscount.org
- Shore, R. & Shore, B. (2009). KIDS COUNT indicator brief: Preventing low birthweight. Baltimore, MD: The Annie E. Casey Foundation.
- ¹² Martin, J., et al. (2013). Births: Final data for 2011. National Vital Statistics Reports, 62(1). Hyattsville, MD: National Center for Health Statistics.

Sources for all other data on health outcomes for RIte Care, Medicaid, private insurance and uninsured in Rhode Island:

Griffin, J. (2013). Health indicator data book: A comparison of access and quality measures for Rhode Islanders <65 years old by health insurance coverage — trends 2001-2011. Cranston, RI: Rhode Island Medicaid Research and Evaluation Project, Rhode Island Department of Human Services.

Griffin, J. (2006). The impact of RIte Care on the health of pregnant women and their newborns 1993-2004. Cranston, RI: Rhode Island Medicaid Research and Evaluation Project, Rhode Island Department of Human Services.