

Women with Delayed Prenatal Care

DEFINITION

Women with delayed prenatal care is the percentage of women receiving prenatal care beginning in the second or third trimester of pregnancy. Data are reported by place of mother's residence, not place of infant's birth.

SIGNIFICANCE

Early prenatal care is an important way to identify and treat health problems as well as influence health behaviors that can affect fetal development, infant health, and maternal health. Women receiving late or no prenatal care are at increased risk of poor birth outcomes, such as having babies who are low birthweight or who die within the first year of life.^{1,2}

Effective prenatal care screens for and intervenes with a range of maternal needs including nutrition, social support, mental health, smoking cessation, substance use, domestic violence, and unmet needs for food and shelter. A prenatal visit is the first step in establishing an infant's medical home and can provide valuable links to other services.^{3,4}

Early prenatal care is especially important for women who face multiple risks for poor birth outcomes, as is ensuring access to health care services before pregnancy. Effective monitoring and treatment of chronic disease,


providing health education, implementing and enhancing Medicaid policies, improving health insurance coverage, and ensuring access to culturally and linguistically competent health providers can improve prenatal care for women of childbearing age.^{5,6}

Barriers to prenatal care include not knowing one is pregnant, not being able to get an appointment or start care when desired, lack of transportation or child care, inability to get time off work, and financial constraints (including lack of insurance or money to pay for desired care).⁷ Rhode Island women with delayed prenatal care are more likely to report their pregnancy was unintended than women who initiated care in the first trimester. Access to contraception, preventative health care services, and the overall health and economic well-being of individuals impact pregnancy intention.^{8,9}

Maternal health before pregnancy (preconception), during pregnancy, and after birth (postpartum) impact health outcomes. Currently, there is a maternal health crisis nationally and in Rhode Island. Beyond that, there are persistent racial and ethnic disparities that disproportionately impact health outcomes for Black, Indigenous, People of Color (BIPOC) women.¹⁰



Women With Delayed or No Prenatal Care by Race/Ethnicity, Rhode Island, 2017-2021



Source: Rhode Island Department of Health, Center for Health Data and Analysis, Maternal and Child Health Database, 2017-2021. * Race categories are non-Hispanic. ^Due to small numbers please interpret percentage with caution.

◆ In Rhode Island between 2017 and 2021, 15.9% of women who gave birth did not begin care until the second or third trimester if at all. Between 2017 and 2021 in Rhode Island, Native Hawaiian or Other Pacific Islander (44.4%), American Indian and Alaska Native (21.7%), Black (21.5%), Hispanic (18.4%), and Asian women (17.9%) were more likely to receive delayed prenatal care than white women (13.2%).¹¹

◆ Between 2017 and 2021 in Rhode Island, women who did not graduate from high school were more likely to receive delayed prenatal care than women with more than a high school education (25.2% compared to 13.0%). Adolescent and teen mothers were more likely to receive delayed prenatal care than older mothers in Rhode Island.¹² About one in five (19.8%) pregnant women in the four core cities received delayed prenatal care compared to 13.4% in the remainder of the state.¹³



Insurance Coverage Improves Access to Prenatal Care

◆ In the U.S. and Rhode Island, women with private insurance have the highest rates of timely prenatal care. Health care before pregnancy is important for maintaining women's reproductive health and ensuring that they can access the reproductive health services they may need to become pregnant, if and when they want to.^{14,15}

◆ Between 2017 and 2021, women with health coverage through RIte Care (Rhode Island's Medicaid managed care program) were much less likely (20.0%) to receive delayed/no prenatal care than women who were uninsured (39.9%). Women with private insurance coverage were the least likely to receive delayed/no prenatal care (11.9%).¹⁶



Racial/Ethnic Disparities in Severe Maternal Morbidity

◆ Nationally, Black women are three times more likely than white women to die of pregnancy-related complications.^{17,18} Racial disparities in maternal mortality span all levels of education, age, income, and insurance status.^{19,20}

◆ Pervasive racial bias and unequal treatment of Black women in the health care system often result in inadequate treatment for pain.^{21,22} This, coupled with stress from racism and racial discrimination, contribute to the unacceptable health outcomes among Black women and their infants.^{23,24}

◆ In Rhode Island, maternal mortality numbers are too small to report. To better measure maternal health during pregnancy and after childbirth, Rhode Island reports the prevalence of severe maternal morbidity. Severe maternal morbidity is defined as unintended outcomes of labor and delivery that result in significant consequences to a woman's health.²⁵

◆ In 2021, the Rhode Island severe maternal morbidity rate was 85 per 10,000 delivery hospitalizations up from 72 per 10,000 in 2020. Black women (112 per 10,000) and Hispanic women (99 per 10,000) had higher rates of maternal morbidity than white women (78 per 10,000) between 2017 and 2021.²⁶

Table 18. Delayed Prenatal Care, Rhode Island, 2017-2021

CITY/TOWN	# BIRTHS	# DELAYED CARE	% DELAYED CARE
Barrington	544	82	15.1
Bristol	660	95	14.4
Burrillville	593	85	14.3
Central Falls	1,457	312	21.4
Charlestown	264	23	8.7
Coventry	1,422	152	10.7
Cranston	3,691	580	15.7
Cumberland	1,603	228	14.2
East Greenwich	535	60	11.2
East Providence	2,147	317	14.8
Exeter	231	22	9.5 ^
Foster	200	27	13.5
Glocester	326	51	15.6
Hopkinton	327	32	9.8
Jamestown	134	11	8.2 ^
Johnston	1,290	192	14.9
Lincoln	858	131	15.3
Little Compton	70	10	14.3 ^
Middletown	786	90	11.5
Narragansett	259	27	10.4
New Shoreham	24	5	*
Newport	1,053	158	15.0
North Kingstown	1,065	117	11.0
North Providence	1,525	231	15.1
North Smithfield	424	75	17.7
Pawtucket	4,196	776	18.5
Portsmouth	655	70	10.7
Providence	11,409	2,306	20.2
Richmond	299	28	9.4
Scituate	420	70	16.7
Smithfield	717	116	16.2
South Kingstown	811	79	9.7
Tiverton	542	70	12.9
Warren	405	64	15.8
Warwick	3,509	417	11.9
West Greenwich	239	29	12.1
West Warwick	1,446	186	12.9
Westerly	887	89	10.0
Woonsocket	2,433	461	18.9
Unknown**	226	32	14.2
Four Core Cities	19,495	3,855	19.8
Remainder of State	29,961	4,019	13.4
Rhode Island	49,682	7,906	15.9

Source of Data for Table/Methodology

Rhode Island Department of Health, Center for Health Data and Analysis, Maternal and Child Health Database, 2017-2021.

The denominator is the total number of live births to Rhode Island residents from 2017-2021.

*The data are statistically unreliable and rates are not reported and should not be calculated.

^The data are statistically unstable and rates or percentages should be interpreted with caution.

**Unknown: Specific city/town information unavailable

Core cities are Central Falls, Pawtucket, Providence, and Woonsocket.

Due to birth certificate changes that began in 2015, comparisons with previous years should be made with caution. Delayed prenatal care is now a calculated variable that is based on the number of visits over 90 days (3 months). "No prenatal care" is not broken out.

References

- Yogman, M., Lavin, A., & Cohen, G. (2018). The prenatal visit. *Pediatrics* 142(1): e20181218.
- U.S. Department of Health & Human Services, Office on Women's Health. (n.d.). *Prenatal care*. Retrieved February 23, 2022, from www.womenshealth.gov
- Hagan, J. F., Shaw, J. S., & Duncan, P. M. (Eds.). (2017). *Bright futures: Guidelines for health supervision of infants, children, and adolescents (4th ed.)*. Elk Grove Village, IL: American Academy of Pediatrics.
- Shore, R. & Shore, B. (2009). *KIDS COUNT indicator brief: Reducing infant mortality*. Baltimore, MD: The Annie E. Casey Foundation.
- Kim, H., Cain, R., & Viner-Brown, S. (2014). *2014 Rhode Island Pregnancy Risk Assessment Monitoring System data book*. Providence, RI: Rhode Island Department of Health.
- Kim, H., Monteiro, K., Cooper, T., Viner-Brown, S., & Weber, A. (2018). *2018 Rhode Island Pregnancy Risk Assessment Monitoring System data book: 3rd edition*. Providence, RI: Rhode Island Department of Health.

(continued on page 181)